

BREAST AUGMENTATION POSTOPERATIVE INSTRUCTIONS & ADVICE

The following after-care information will help make your surgery safe and uneventful. It is important that you understand it all. If you have any questions, please ask me or my staff. In some cases there will be variations to the instructions, and these will be discussed with you.

Dressing

After a breast augmentation, you will find steri strips and white tape on the scar line. This tape should stay on for up to 2 weeks. After your first post-operative appointment, you can start showering normally over the tape. The tapes will come off on their own in the shower but should be removed if still on at 2 weeks.

Surgical Bra

You will wake up with a surgical bra on. This bra provides support and stabilizes the position of the implants. The dressing and bra should stay on and untouched until your first post-op appointment. After this, you can remove and replace the bra as you wish but it should be worn day and night, as much as possible for 4 weeks. Please note that most of the “seams” of a surgical bra are on the outside of the bra to avoid irritation of the skin.

Pain Pills & Antibiotics

Most patients are pleasantly surprised by how limited their postoperative discomfort is. You have been prescribed Tylenol extra strength, Celebrex (Celecoxib) which is an anti-inflammatory and Hydromorphone which is a strong narcotic pain killer. The Tylenol should be taken on schedule, every 6 hours. For the first 3 days, the Celebrex is to be taken every 12 hours. The Hydromorphone is to be taken as needed but not more often than every 6 hours.

This combination of anti-inflammatory (Celebrex) and analgesic (Tylenol extra strength or stronger pain pill) is now recognized as the most effective way of treating postoperative discomfort. Both can be stopped when the discomfort subsides. Please note that all pain pills cause constipation, and if you are inclined to becoming constipated, you should consider the use of a stool softener such as Colace 200 mg twice a day, and a laxative such as Milk of Magnesia, 15 to 30 cc at nighttime. Drinking copious amounts of fluids and eating fruit also reduces constipation.

Antibiotics are prescribed routinely to minimize the chance of infection and the full course of antibiotics should be taken.

Implant Massage

At your first post-operative visit you will be shown how to massage the implant. Starting 1 week after surgery, you will massage the implant daily to move it inwards and upwards within the breast pocket. This maneuver should be performed 5 times in each direction for each breast. This should be repeated twice daily for 2 months then once per day indefinitely.

Eating After Surgery

Patients may experience nausea after a general anesthetic. It is therefore wise to drink clear fluids (apple juice, black tea, flat pop) until the stomach feels settled. Then try eating clear soup and a few crackers. A normal diet is usually possible the next day. On rare occasion, a patient may feel nausea for a few days after surgery. Gravol tablets may then be helpful.

Hematoma

On rare occasion, a hematoma can develop. This is due to continued bleeding underneath the skin. It presents itself as an enlarging, painful, firm swelling of the entire breast, typically in the first 24 to 48 hours after surgery. Small differences in size and swelling are normal. A hematoma is quite obviously a major difference in size. It may require surgery and an urgent evaluation by me or one of the emergency room physicians at North York General Hospital is necessary.

Bruising

Bruising is the typical "black and blue" appearance to tissue after any surgery. There is a great variation in how much patients bruise, some bruising very little, some more. As noted in your preoperative instructions, Aspirin, blood thinners, anti-inflammatories, and vitamins can contribute to bruising and should be stopped for two weeks before surgery. These medications can be resumed the day after surgery. The bruising typically subsides within two weeks, but in some patients can last longer. As it dissolves, and becomes more yellowish in colour, it may appear to be "spreading". This is normal and is no cause for alarm. It is common to have more bruising and swelling on one side than the other. This will even out with time.

Arnica for Bruising

Arnica has been shown to help decrease bruising and reduce the length that bruising is visible.

It is recommended to use a combination of oral pellets and topical gel but if you would only like to use one or the other, the pills are likely more effective. The pills should be started two days before your procedure and continued for 3-5 days after your procedure. The gel can be started as soon as the massage process of administering the gel is tolerable and can be continued until the bruising has subsided. Storing the gel in the fridge will also provide a cooling and soothing effect to the bruised area.

DO NOT take off the dressing and DO NOT apply on the actual incision until completely healed. Apply around procedure area to minimize swelling and bruising. Please DO NOT use Arnica if you are allergic to the daisy flower.

Early Fever

A low grade fever (up to 37.8°C) is not uncommon the day after surgery. It usually occurs after a general anesthetic and results from small collections of mucous in the lungs. It is therefore important after a general anesthetic to take frequent deep breaths, and to cough deeply. The coughing clears the lungs and corrects the low grade fever. Failure to clear the lungs adequately can on rare occasion lead to a lung infection.

Wound Infection

Wound infections fortunately are uncommon. Minor leakage of clear yellowish fluid from stitches may occur and persist until the stitches have dissolved. This fluid is not an infection and is managed with soap and water washes two to three times a day, followed by the application of Polysporin ointment and a light dressing. If the drainage from the wound becomes “soupy” and smells, or is associated with increasing redness, pain and swelling, then a deeper infection may be occurring. Do not hesitate to phone the office or go to the emergency room department if this happens. You will need to be seen as you may need to have antibiotics.

Sleeping Position

The incision lines are very secure, and you can sleep on your back or on either side immediately after surgery. It is suggested that you do not sleep on your stomach for 4 weeks after surgery.

Scar Cream

Once the tapes are off the scar line, a scar gel should be used. The gel is first massaged onto the incision lines. Additional scar gel is then placed on the scar lines, allowed to dry and covered with Paper tape. This process should be repeated every other day. You can shower over the tape on the intervening days. The scar cream I recommend is available in our office.

Polysporin Ointment

Occasionally, there are small areas of drainage along the incision line. These little areas of delayed healing can open up slightly and a clear yellowish fluid may drain out. This is not infection. These areas should be washed with soap and water twice a day, followed by Polysporin ointment and gauze. Scar cream should be avoided in these areas until the wounds have completely healed, whereupon the scar cream can also be applied. While these small areas are healing, scar cream is still used on the remaining incision lines.

Wound Massage

As soon as the scar cream is being applied, 10 to 14 days after surgery, gentle massage of the incision lines should also be carried out. After another week, this gentle massage can become quite firm and vigorous, with deep rotary motions along the incision line. This firm massage helps to desensitize the scars, making them less painful, and also reduces the normal scar thickness and lumpiness which is seen in a scar after surgery.

Scar Redness

All scars initially are reddish and thickened. This is part of the normal healing process and eventually will smoothen out. It takes 9 to 12 months for the scars to undergo a full maturation process whereby they become flat, light in colour, and somewhat more spread.

Tanning

Fresh scars and areas that have been bruised should not be tanned for at least 6 months after surgery. Early tanning can cause permanent hyperpigmentation of the scar and of the surrounding areas. Normal tanning can resume after four months, but please remember that tanning does cause premature skin aging and skin cancers.

Silicone Sheeting

Some studies have suggested that silicone sheeting, available in most pharmacies and on the internet, can help the quality of the scar (Cica-care or Scar-away). To be effective, this silicone sheeting needs to be on the skin day and night for at least six months. It is not recommended for routine usage as it appears to have little value when scars are healing normally. I do recommend it in wounds that are showing evidence of prolonged redness or abnormal scar maturation. This is in less than 5% of patients.

Mobility

As a routine, it is important to be up and walking the day of surgery, and prolonged periods of bed rest are discouraged. Lying in bed can increase the rare but ever-present risk of developing blood clots in the calves which in turn can cause pulmonary emboli. It is good to go for little walks, and to have frequent naps if you are feeling tired. For the first two weeks there should not be any excessive stretching on the wound, but even the day after surgery, you can be gently raising your arms as needed. Within 4 weeks, full exercise activities can be resumed.

Driving a Car

You must not be driving a car for at least 24 hours after a sedative or general anesthetic, nor if you feel drowsy for any reason. Do not drive a motor vehicle if you are having any pain, as the pain may cause you to jerk the steering wheel and lose control. You also must avoid driving a car if you are in any way restricted in your mobility as this may compromise safe driving.

Fatigue and “Feeling Down”

While the results of plastic surgery are typically gratifying and uplifting, it is not uncommon for patients to have “down” times. Early on there is the residual effect of the anesthetic, coupled with discomfort, swelling and bruising at the surgical site. This is followed by the typically slow maturation of the scar itself. These frustrations all play together on the psyche, making one question the reason for doing the surgery in the first place. You are not alone with these thoughts, and they are entirely normal. Rest assured that well over 90% of patients are pleased with the results of their surgery, and the apprehensions and “down” times are eventually replaced by feelings of satisfaction and confidence.

Questions/Problems

Every effort is made to provide you with a surgical experience that is safe and as comfortable as possible. Any suggestions you have that might improve the experience are much appreciated.

If you have any questions, concerns, or problems, please call the office at 416-447-6176. If there is an urgent situation, and we are not immediately available, please go to the emergency room department at North York General Hospital, or your closest emergency room department.

We hope you have a speedy recovery from your surgery.

Sincerely,



Ron B. Somogyi, M.D.



Waqqas Jalil, M.D.