

LIPOSUCTION/FAT INJECTION

The following after-care information will help make your surgery safe and uneventful. It is important that you understand it all. If you have any questions, please ask me or my staff. In some cases there will be variations to the instructions, and these will be discussed with you.

Wound Care

After surgery you will have small see-through Op-Site dressings on the tiny liposuction incision points. Please keep all bandages dry until your first post-operative visit. The gauze underneath the plastic dressing may become saturated with pink fluid/blood – this is normal. We will remove these op-site dressings and you will see small steri strips over the scars, you can shower normally over these and they will come off or should be removed by 2 weeks post-operatively. The stitches, if visible, will fall out by themselves.

Once the tapes are off, a scar cream should be massaged onto the incisions twice a day for two to three months. If any area of drainage remains, Polysporin ointment, rather than the scar cream, should be applied on those areas until the drainage has stopped. The scar cream is then also applied there. The scar cream I recommend is available in our office.

Compression Garment

You will wake up with a compression garment in place. This garment reduces bruising and swelling and allows the skin to redrape over the new body contour. It should be worn for three weeks night and day, and then for three weeks only during the day. These garments can be washed in a washing machine. They should be hung dry or placed in the tumble (rather than heat) cycle of the dryer.

Pain Pills

Most patients are pleasantly surprised by how limited their postoperative discomfort is. You have been prescribed Tylenol extra strength, Celebrex (Celecoxib) which is an anti-inflammatory and Hydromorphone which is a strong narcotic pain killer. The Tylenol should be taken on schedule, every 6 hours. For the first 3 days, the Celebrex is to be taken every 12 hours. The Hydromorphone is to be taken as needed but not more often than every 6 hours.

This combination of anti-inflammatory (Celebrex) and analgesic (Tylenol extra strength or stronger pain pill) is now recognized as the most effective way of treating postoperative discomfort. Both can be stopped when the discomfort subsides. Please note that all pain pills cause constipation, and if you are inclined to becoming constipated, you should consider the use of a stool softener such as Colace 200 mg twice a day, and a laxative such as Milk of Magnesia, 15 to 30 cc at nighttime. Drinking copious amounts of fluids and eating fruit also reduces constipation.

Bleeding, Bruising, Swelling

Minor bleeding from the incision lines is normal after surgery, and should subside within 24 hours. Bruising is the typical "black and blue" appearance to tissue after any surgery. There is a great variation in how much patients bruise, some bruising very little, some more. As noted in your preoperative instructions, Aspirin, blood thinners, anti-inflammatories, and vitamins can contribute to bruising and should be stopped for two weeks before surgery. These medications can be resumed the day after surgery. The bruising typically subsides within two weeks, but in some patients can last longer. As it dissolves, and becomes more yellowish in colour it may appear to be "spreading". This is normal and is no cause for alarm. It is common to have more bruising and swelling on one side than the other. This will even out with time.

Arnica for Bruising

Arnica has been shown to help decrease bruising and reduce the length that bruising is visible.

It is recommended to use a combination of oral pellets and topical gel but if you would only like to use one or the other, the pills are likely more effective. The pills should be started two days before your procedure and continued for 3-5 days after your procedure. The gel can be started as soon as the massage process of administering the gel is tolerable and can be continued until the bruising has subsided. Storing the gel in the fridge will also provide a cooling and soothing effect to the bruised area.

DO NOT take off the dressing and DO NOT apply on the actual incision until completely healed. Apply around procedure area to minimize swelling and bruising. Please DO NOT use Arnica if you are allergic to the daisy flower.

Early Fever

A low grade fever (up to 37.8°C) is not uncommon the day after surgery. It usually occurs after a general anesthetic and results from small collections of mucous in the lungs. It is therefore important after a general anesthetic to take frequent deep breaths, and to cough deeply. The coughing clears the lungs and corrects the low grade fever. Failure to clear the lungs adequately can on rare occasion lead to a lung infection.

Wound Infection

Wound infections fortunately are uncommon. Minor leakage of clear yellowish fluid from stitches may occur and persist until the stitches are out. This fluid is not an infection, and is managed with soap and water washes two to three times a day, followed by the application of Polysporin ointment and a light dressing.

If the drainage from the wound becomes "soupy" and smells, or is associated with increasing redness, pain and swelling, then a deeper infection may be occurring. Do not hesitate to phone the office if this happens. You will need to be seen as you may need to have antibiotics.

Wound Massage

As soon as the scar cream is being applied, 10 to 14 days after surgery, gentle massage of the incision lines should also be carried out. After another week, this gentle massage can become quite firm and vigorous, with deep rotary motions along the incision line. This firm massage helps to desensitize the scars, making them less painful, and also reduces the normal scar thickness and lumpiness which is seen in a scar after surgery.

Massage of the Liposuction Area

One week after your surgery, you can start to gently massage the areas that have been contoured. You can use a circular motion and try to evenly cover the entire area. After 2 weeks this massage can be more vigorous. A good moisturizer or Bio-oil can be used for this. Spend at least 5 minutes once a day doing this for 6 weeks.

Scar Redness

All scars initially are reddish and thickened. This is part of the normal healing process and eventually will smoothen out. It takes 9 to 12 months for the scars to undergo a full maturation process whereby they become flat, light in colour, and somewhat more spread.

Tissue Swelling

While 80% of the swelling will be gone in three weeks, the last 20% of swelling can also take 9 to 12 months to completely subside.

Tanning

Fresh scars and areas that have been bruised should not be tanned for at least 6 months after surgery. Early tanning can cause permanent hyperpigmentation of the scar and of the surrounding areas. Normal tanning can resume after four months, but please remember that tanning does cause premature skin aging and skin cancers.

Mobility

As a routine, it is important to be up and walking the day of surgery, and prolonged periods of bed rest are discouraged. Lying in bed can increase the rare but ever-present risk of developing blood clots in the calves which in turn can cause pulmonary emboli. It is good to go for little walks, and to have frequent naps if you are feeling tired. For the first two weeks there should not be an heavy lifting or exercise that makes your heart beat fast. More bleeding could result. After two weeks, full activities can be resumed. You can return to full exercise by 3 weeks but, note, you may see temporary increased swelling after periods of increased activity for the first few months.

Driving a Car

You must not be driving a car for at least 24 hours after a sedative or a general anesthetic, nor if you feel drowsy for any reason. Do not drive a motor vehicle if you are having any pain, as the pain may cause you to jerk the steering wheel and lose control. You also must avoid driving a car if you are in any way restricted in your mobility or your vision as this may compromise safe driving.

Fatigue and “Feeling Down”

While the results of plastic surgery are typically gratifying and uplifting, it is not uncommon for patients to have “down” times. Early on there is the residual effect of the anesthetic, coupled with discomfort, swelling and bruising at the surgical site. This is followed by the typically slow maturation of the scar itself. These frustrations

all play together on the psyche, making one question the reason for doing the surgery in the first place. You are not alone with these thoughts, and they are entirely normal. Rest assured that well over 90% of patients are pleased with the results of their surgery, and the apprehensions and “down” times are eventually replaced by feelings of satisfaction and confidence.

Questions/Problems

Every effort is made to provide you with a surgical experience that is safe and as comfortable as possible. Any suggestions you have that might improve the experience are much appreciated.

If you have any questions, concerns, or problems, please call the office at 416-447-6176. If there is an urgent situation, and we are not immediately available, please go to the emergency room department at North York General Hospital, or your closest emergency room department.

We hope you have a speedy recovery from your surgery.

Sincerely,



Ron B. Somogyi, M.D.



Waqqas Jalil, M.D.